

# Surgifort® Human Milk Fortifier (Human, Pasteurized)

To be used under the supervision of a physician in the hospital. Intended for term infants with Gastroschisis following surgical repair.

## Important Reminders

- Store Surgifort fortifier at  $-20^{\circ}\text{C}$  or colder until ready to thaw for use
- Always refer to the Supplemental Product Information (SPI) for Surgifort fortifier when preparing the feeding solution (available online at [Prolacta.com](http://Prolacta.com))
- Human milk varies in protein content with lactation stage. Protein intake varies as target caloric concentration changes. Nutrition monitoring is required.
- Do not add medications, supplements, or non-human milk-based modulars directly to the Surgifort fortifier and human milk feeding solution
- Any transition of enteral feeding source can result in gastrointestinal complications such as intolerance or more serious issues. Consider a phased transition over multiple days to allow the infant to acclimate to the new food source.



Product images shown are for illustration purposes only

## Thawing

Under no circumstances should the product be defrosted or warmed in a microwave.

- Remove product container from freezer and label with date and time
- Place unopened (frozen) product container in refrigerator ( $2^{\circ}\text{C}$  to  $8^{\circ}\text{C}$ ) for 2–5 hours
- Swirl gently to detect ice in the product container; if ice is still present, return to the refrigerator for additional thaw time; repeat until no ice is detected
- Once the thawing process begins, administer within 48 hours; discard any unused portion after mixing
- Do not refreeze; keep refrigerated until used


## Preparation

Always maintain aseptic technique when preparing and handling human milk. This is a concentrated liquid, **DO NOT ADD** human milk to the product container. **DO NOT ADD WATER.**

Pour required quantity of Surgifort fortifier into a container holding the required volume of human milk.

### Nutrient Values Of Surgifort Feeding Solution When Mixed With Term Human Milk†

Mixing Ratio Surgifort fortifier:Human milk	Surgifort Fortifier	Term Human Milk	Total	Calories per 100 mL	Protein per 100 mL
3:7 to achieve <b>26 kcal/fl oz</b>	30 mL	70 mL	100 mL	88.5 kcal	1.8 g
2:3 to achieve <b>28 kcal/fl oz</b>	40 mL	60 mL	100 mL	95.4 kcal	2.1 g



For complete information, call 1.888.776.5228 or visit [Surgifort.com](http://Surgifort.com)

†American Academy of Pediatrics Committee on Nutrition. Appendix A. In: Kleinman RE, Greer FR, eds. *Pediatric Nutrition*. 8th ed. Itasca, IL: American Academy of Pediatrics; 2019:1505-1508.